



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Haptoglobin and hemoglobin in subarachnoid hemorrhage: A tale of 2 globins

Loch Macdonald, R ; Katan, Mira

DOI: <https://doi.org/10.1212/WNL.0000000000007399>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-176664>

Journal Article

Published Version

Originally published at:

Loch Macdonald, R; Katan, Mira (2019). Haptoglobin and hemoglobin in subarachnoid hemorrhage: A tale of 2 globins. *Neurology*, 92(18):831-832.

DOI: <https://doi.org/10.1212/WNL.0000000000007399>

Haptoglobin and hemoglobin in subarachnoid hemorrhage

A tale of 2 globins

R. Loch Macdonald, MD, PhD, and Mira Katan, MD, MS

Neurology® 2019;92:831-832. doi:10.1212/WNL.00000000000007399

Correspondence

Dr. Macdonald
rlochmacdonald@gmail.com

Although the incidence is declining, subarachnoid hemorrhage (SAH) continues to exact a substantial toll on society, with an estimated 20,000 aneurysmal SAHs (aSAHs) a year in the United States.¹ Outcomes have improved, but the mortality remains ≈35%, and half of the survivors cannot return to their previous level of functioning.² The most important prognostic factors for outcome are the neurologic condition of the patient on admission to hospital, the patient's age, and preexisting hypertension.³ aSAH has a unique biphasic course, with initial or early brain injury (reflected in the neurologic condition) and a delayed phase of brain injury called delayed cerebral ischemia (DCI). Subarachnoid blood and specifically the erythrocytes and their main content, hemoglobin, appear to mediate DCI in humans. Extravascular free hemoglobin and iron released from it, whether into the CSF or the brain, are highly inflammatory and cytotoxic oxidants.⁴ The body has developed ways to mitigate this, including synthesis of haptoglobin, which binds free hemoglobin, abrogating its toxicity. Therefore, clearing away the hemoglobin more quickly or preventing its deleterious effects by binding hemoglobin may decrease brain injury and improve patient outcomes from SAH.

Humans have 2 haptoglobin alleles, haptoglobin 1 and 2, that differ in their scavenging ability.⁴ Of the 3 possible genotypes, the haptoglobin 2-2 genotype may have reduced scavenging and has associations with more severe phenotypes of diseases such as diabetes mellitus, cardiovascular disease, and renal failure.⁴ Patients with haptoglobin 2-2 appear more likely to develop elevated transcranial Doppler intracranial arterial flow velocities, an indicator of vasospasm of the intracranial arteries after SAH,⁵ although subsequently published data conflict on the role of haptoglobin genotype plays in SAH outcome.

In this issue of *Neurology*®, Gaastra et al.⁶ performed an important study by collecting a large dataset that includes haptoglobin genotypes and outcome after aSAH. The fact that the authors collected 6 unpublished datasets along with 5 published ones demonstrates important collaborations with the authors.⁶ The rigorously performed individual patient-level meta-analysis of these 939 patients, however, showed no evidence for any effect of haptoglobin genotype on the prespecified outcomes, including functional outcome, short-term complications of aSAH such as cerebral infarction, DCI, angiographic vasospasm, and transcranial Doppler flow velocities.

What reasons underlie this lack of association? Haptoglobin genotype may not influence the outcome. On the other hand, aSAH is a complex disease with many different patient-, disease-, and treatment-related factors contributing to patient outcome, again requiring large datasets to detect small prognostic effects. For example, treatment intensity may obviate the effects of haptoglobin genotype. Moreover, the modified Rankin Scale or Glasgow Outcome Scale scores dichotomized into favorable and unfavorable outcome—a rather blunt outcome assessment—may not have adequate sensitivity to detect small changes in patient outcome. In summary, this study demonstrates that if haptoglobin genotype does

RELATED ARTICLE

Haptoglobin genotype and aneurysmal subarachnoid hemorrhage: Individual patient data analysis

Page 844

From the Division of Neurosurgery (R.L.M.), St. Michael's Hospital, Labatt Family Centre of Excellence in Brain Injury and Trauma Research, Keenan Research Centre for Biomedical Research and Li Ka Shing Knowledge Institute, St. Michael's Hospital, Department of Surgery, University of Toronto, Ontario, Canada; and Stroke Center (M.K.), Department of Neurology, University Hospital of Zurich, Switzerland.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

have an effect on outcome, the effect size would be small and most probably not clinically relevant.

Roughly 4.5×10^{19} molecules of hemoglobin are released from the typical aSAH volume of ≈ 35 mL into the ≈ 50 -mL subarachnoid space, with each hemolyzing erythrocyte releasing 250 million molecules.^{7,8} As an acute-phase reactant, the concentration of haptoglobin in the CSF, normally ≈ 0.06 mg/dL, would increase but would likely still be overwhelmed. Little is known about the concentrations of haptoglobin or of the different proteins resulting from the different genotypes in the CSF and brain after SAH; this remains an important area for additional work.

This study clearly demonstrates the large numbers of patients required to move the field along and highlights the need for national and international collaborations. Finally, as the authors mention, different studies collect different variables, use different definitions, and follow up patients for varying times. Common data elements are going to be key in the efforts to study large numbers of patients with aSAH.

Study funding

No targeted funding reported.

Disclosure

R. L. Macdonald receives grant support from the Brain Aneurysm Foundation and Genome Canada. M. Katan receives grant support from the Swiss National Science Foundation, Swiss Heart Foundation, and University Hospital of Zurich, Switzerland. Go to Neurology.org/N for full disclosures.

References

1. Etminan N, Chang HS, Hackenberg K, et al. Worldwide incidence of aneurysmal subarachnoid hemorrhage according to region, time period, blood pressure, and smoking prevalence in the population: a systematic review and meta-analysis. *JAMA Neurol* Epub 2019 Jan 19.
2. Al-Khindi T, Macdonald RL, Schweizer TA. Cognitive and functional outcome after aneurysmal subarachnoid hemorrhage. *Stroke* 2010;41:e519–e536.
3. Jaja BNR, Saposnik G, Lingsma HF, et al. Development and validation of outcome prediction models for aneurysmal subarachnoid haemorrhage: the SAHIT multinational cohort study. *BMJ* 2018;360:j5745.
4. Bulters D, Gaastra B, Zolnourian A, et al. Haemoglobin scavenging in intracranial bleeding: biology and clinical implications. *Nat Rev Neurol* 2018;14:416–432.
5. Borsody M, Burke A, Coplin W, Miller-Lotan R, Levy A. Haptoglobin and the development of cerebral artery vasospasm after subarachnoid hemorrhage. *Neurology* 2006;66:634–640.
6. Gaastra B, Ren D, Alexander S, et al. Haptoglobin genotype and aneurysmal subarachnoid hemorrhage: individual patient data analysis. *Neurology* 2019;92:e2150–e2164.
7. Whitmore RG, Grant RA, Leroux P, El-Falaki O, Stein SC. How large is the typical subarachnoid hemorrhage? A review of current neurosurgical knowledge. *World Neurosurg* 2012;77:686–697.
8. Wintrobe MM, Greer JP. *Wintrobe's Clinical Hematology*, 11th ed. Philadelphia: Lippincott Williams & Wilkins; 2004.